

Routine Publicly Funded Immunization Schedule for Ontario (Children Starting Immunization in Infancy*)

*Additional doses may be needed for high risk individuals, or for Catch-Up Schedules. Refer to *Publicly Funded Immunization Schedules for Ontario* (2016 or as current) at www.health.gov.on.ca/en/pro/programs/immunization/schedule.aspx
For more up-to-date information, contact Hastings Prince Edward Public Health at 613-966-5500 or 1-800-267-2803 / TTY 711 or 1-800-267-6511 / www.hpepublichealth.ca

Immunizing Agent / OHIP Billing Code	2 Mos	4 Mos	6 Mos	12 Mos	15 Mos	18 Mos	4-6 Yrs	Gr 7-8	Gr 7-12	Gr 7	9-26 Yrs	14-16 Yrs	Every 10 Yrs	65+ Yrs
Diphtheria, tetanus, pertussis, polio, Hib Pediactel / Pentacel G841	✓1	✓1	✓1			✓1								
Pneumococcal conjugate 13 Prenar 13 G846	✓2	✓2		✓2										
Rotavirus RotaTeq (3 doses) / Rotarix (2 doses) N/A	✓3	✓3	✓3											
Meningococcal C conjugate Menjugate / NeisVac-C / Meningitec G844				✓4										
Measles, mumps, rubella (MMR) MMRII / Priorix G845				✓5										
Varicella Varivax III / Varilrix G848					✓6									
Measles, mumps, rubella, varicella (MMRV) Priorix-Tetra / Pro Quad G538							✓7							
Tetanus, diphtheria, pertussis, polio Adacel-Polio / Boostrix-Polio G538							✓8							
Hepatitis B Recombivax HB / Engerix B G842								✓9						
Meningococcal conjugate ACYW-135 Menactra G538									✓10					
Human papillomavirus (HPV9) Gardasil G843										✓11	✓11			
Tetanus, diphtheria, pertussis Adacel / Boostrix G847												✓12	✓12	
Tetanus, diphtheria G538													✓13	
Pneumococcal polysaccharide 23 Pneumo 23 G538														✓14
Shingles (Zostavax) G538														✓15
Influenza G590										✓16				

1	Pediacel or Pentacel (DTap-IPV-Hib) is given at 2, 4, 6 and 18 mos of age to complete the primary series.
2	Prevnar 13: Healthy children receive Prevnar 13 at 2, 4, and 12 months of age.
3	RotaTeq: Given orally, usually at 2, 4 and 6 months of age. May be administered as early as 6 weeks and up to 32 weeks. First dose should be before 15 weeks of age; 2nd dose 4 to 10 weeks after first; 3rd dose 4 to 10 weeks after the second. Rotarix: Given orally at 2 and 4 months of age. First dose should be given between 6 weeks of age and before 15 weeks of age; the second dose is given before 25 weeks of age. Doses must be at least 4 weeks apart.
4	Meningococcal-C is given at 12 mos of age (or up to grade 6 if missed).
5	MMR should be given on or after the first birthday. A second dose is given at 4 to 6 years of age combined with varicella (MMRV).
6	Varicella: a two-dose schedule of varicella is recommended. The first dose is given at 15 months as single varicella; the second dose is given at 4 to 6 years of age in a vaccine combined with MMR (MMRV).
7	MMRV is recommended for the second doses of MMR and varicella, and is given at 4 to 6 years of age. In Ontario, it is only recommended from 4 to 12 years of age inclusive.
8	Tetanus, diphtheria, pertussis, polio (Tdap-IPV) is given as the 5th (booster) dose to 4 to 6 year olds to complete their primary childhood series. [Note: in Sept 2017, Pediacel replaced Quadracel/Infanrix-IPV for children less than 7 yrs of age who have never been immunized and are completing their primary series.]
9	Hepatitis B is offered free to all Grade 7 & 8 students under the School-Based Program. It is a two-dose series for those 11-15 yrs of age inclusive (at 0 and 4-6 months, depending on the product) and a 3-dose series for anyone outside this age range (at 0, 1 and 6 mos).
10	Meningococcal quadrivalent conjugate ACYW-135 is free for all Grade 7 to 12 students under the School-Based Program.
11	HPV9 is offered free to all Grade 7 students under the School-Based Program (or as current). MSM (men who have sex with men) aged 9 to 26 years are also eligible under the High-Risk criteria.
12	Tetanus, diphtheria, and pertussis (Tdap) is recommended for 14 to 16 year olds. Also publicly-funded as a one-time dose for adults who have never received an acellular pertussis-containing vaccine.
13	Tetanus, diphtheria (Td) boosters are recommended every 10 years for adults. [Note: Tetanus, diphtheria, and pertussis (Tdap) is publicly funded as a one-time dose for adults, and should be given to those who have never received an acellular pertussis-containing vaccine.]
14	Pneumococcal polysaccharide 23. One life-time dose is recommended for people 65 years of age and older, people at high risk for pneumococcal disease, and residents of long-term care facilities. Certain high-risk individuals require a 2 nd dose 5 years after the first.
15	Shingles: Publicly-funded for individuals 65-70 years of age inclusive.
16	Influenza: One dose every year in the fall. Children 6 months to less than 9 years of age who have never been previously immunized with a seasonal influenza vaccine should receive 2 doses at least 4 weeks apart, thereafter one dose every year.

We are committed to providing accessible publications, programs and services to all. For assistance please call 613-966-5500; TTY: 711; or email accessibility@hpeph.ca. For more information, please visit www.hpepublichealth.ca.