

# The Social Determinants of Health

A Snapshot of Hastings and Prince Edward Counties August 2017

## INTRODUCTION

## TO THE SOCIAL DETERMINANTS OF HEALTH


The social determinants of health are a group of non-biological factors that influence one's health. They are a major cause of health inequity within a population, making some groups of people healthier than others. The Ontario Public Health Standards list a set of determinants which include: income, education, and employment, among others. Addressing these determinants and their effects on health is a key role of public health units in Ontario. This is an important step in improving the health of not only the most vulnerable people, but of the entire local population.

### The social determinants of health are:



This report illustrates the state of these determinants in Hastings and Prince Edward Counties (HPE), with brief explanations of how they may affect one's health. The information contained in this report is intended to inform the community about the state of social determinants in Hastings and Prince Edward Counties, in the hopes that the health inequities that currently exist can be mitigated through partnerships and collaboration.


In this report, maps are used to illustrate how regions within HPE compare to provincial averages for various social determinants of health. This is displayed by dividing the municipal rate by the provincial rate, otherwise known as a rate ratio. For example, a value of 1.4 indicates that the municipal rate is 1.4 times that of Ontario, making it much higher. A value of 0.75 means that the municipal rate is 0.75 times that of Ontario, making it much lower. Values close to 1 mean that the municipal rate and that of Ontario are similar. Maps are coloured based on these rates to give an overview of how the different municipalities fare for each determinant, per the legend.



# A DEMOGRAPHIC PROFILE

The Hastings and Prince Edward Counties public health region<sup>1</sup>:

- is an urban rural mix
- has a higher proportion of Indigenous people
- has a lower proportion of males
- has had slow population growth



of the population in HPE is 60 years of age or older. (Census 2016)



individuals in HPE earn above \$40,000 annually, while 1 in 3 people in Ontario make above the same threshold. (NHS 2011)



The majority of the population in HPE live in **2 person** households, and in single-detached homes. (Census 2016)

Much of the data in this report comes from the Canadian Community Health Survey (CCHS). It is a survey administered by Statistics Canada that collects detailed, self-reported health information.

In addition to the CCHS, data in this report also comes from the National Household Survey (NHS). The NHS replaced the Canadian long-form census in 2011. Due to its voluntary nature, Tyendinaga Township, Marmora, Wollaston and Deseronto had large non-response rates, making the results non-reportable.




# INCOME AND SOCIAL STATUS

Public perceptions about the social determinants of health in two Ontario public health units show that income is not considered to be an important determinant of health.<sup>2,3</sup> However, there is overwhelming evidence that income and social status are key determinants of how healthy a person will be.<sup>4,5</sup> How income and social status affect health is complex and implicates many other determinants of health.


## % of Low-Income Households in HPE Compared to Ontario

**Ontario rate: 13.9% | HPE rate: 14.8%**

In Figure 1, the proportion of households in HPE that are below the “after-tax low-income measure” compared to that of Ontario is shown (see NHS 2011 for more information). Many of the regions in HPE have high rates of low-income households.



## Perceived Healthiness in Ontario by Income



**Figure 2:** Percentage of respondents in Ontario who rated their health as “very good” or “excellent”, by the respondent’s income quintile. Source: CCHS 2015

The poorest people in Ontario feel the least healthy (Figure 2). Many people living in HPE are found in the lower quintiles of this graph; this implies many health challenges due to income exist in HPE.

Those with less money tend to have poorer childhoods, education, employment, working conditions, housing and food security. These factors all contribute to poor health in different ways.

Improvements in perceived health are seen at every step up in income level. This is a common feature in health equity studies, and is called a **health gradient**.

**Figure 1:** Proportion of residents in HPE that are below the after-tax low-income measure, compared to Ontario. Source: NHS 2011


## EDUCATION AND LITERACY

Lower levels of education are associated with poorer health<sup>4</sup> There are many ways that differences in education may lead to differences in health; for example, higher education may lead to higher income and better health literacy.<sup>6</sup> It can also work in reverse; poor health may lead to lower education. This can make it hard for one to better their health, as they may lack opportunities to increase their income or health literacy.<sup>7</sup>


### % Without Secondary School Completion in HPE Compared to Ontario

**Ontario rate: 10.9% | HPE rate: 15.4%**

HPE has a higher proportion of people without secondary school completion when compared to Ontario (Figure 3). This means there is a larger population within HPE that is at risk of poor health due to lower education.



### Smoking and Obesity in Ontario by Education



**Figure 4:** Proportion of residents in Ontario that reported smoking daily, and proportion of residents in Ontario that reported having a BMI score in the obese range, by highest level of education attained. Source: CCHS 2015

**Figure 3:** Proportion of residents in HPE aged 25-64 years without secondary school completion compared to the Ontario rate. Source: NHS 2011

Being health literate means knowing what's good or bad for one's health. Someone who is not well informed may partake in poor health behaviours. This includes things like smoking, and poor eating habits<sup>8</sup> This notion is supported by provincial data which shows that people with less education are also more likely to smoke and be obese (Figure 4). Other issues that stem from low education include higher unemployment rates and riskier employment such as being a truck driver or construction worker.<sup>9</sup>


# EMPLOYMENT AND WORKING CONDITIONS

Being unemployed, underemployed, or working in stressful/unsafe environments can lead to poorer health. Employment provides a sense of identity and purpose; it also provides opportunities for growth and social activity. Unemployed people have reduced life expectancy and suffer more health problems than employed people.<sup>4,8</sup> Full-time employment also usually includes benefits that will allow individuals to seek and obtain services that will enhance their health, such as dental coverage and paid time off work to seek treatments.

## % Unemployed in HPE Compared to Ontario

**Ontario rate: 8.27% | HPE rate: 8.1%**

The unemployment rate for those aged 15+ in HPE is similar to the Ontario rate. Despite this, certain regions within HPE such as Tudor and Cashel, and Faraday have significantly higher levels of unemployment, putting them at a higher risk for poor health.



Recent data from the 2016 Labour Force Survey, a monthly snapshot of the Canadian labour market, shows that for youth aged 15–24, unemployment in HPE is higher than that of Ontario (17% vs. 14%). This indicates that this age group in particular is at a higher risk of health concerns due to unemployment.

There are 3 main factors thought to be linking unemployment to poor health<sup>4</sup>:



### Material deprivation

When one is unemployed they will have a significantly reduced income, and will also no longer have access to benefits provided by employers.



### Mental health concerns

Particularly if one has just lost their job, feelings of inadequacy, a disruption of daily routine and increased anxiety are just some of the issues one could struggle with when unemployed.



### Unhealthy behaviours

Unhealthy coping behaviours such as smoking or excessive drinking can be more serious issues in the unemployed population.

**Figure 5:** Unemployment rates of residents aged 15+ in HPE compared to the unemployment rate of Ontario. Source: NHS 2011


## SOCIAL SUPPORT NETWORKS

Having people to rely on and provide comfort when one is going through hard times is important to mental and physical health. The social support network determinant of health has to do with personal relationships one has with close friends and family. These contacts may provide a boost to health especially when one is in stressful situations or facing major life changes.

### Social Deprivation in HPE Compared to Ontario

Social deprivation can be measured many different ways. Three common indicators are<sup>10</sup>:

- People living alone
- People that are separated, divorced or widowed
- Number of single parent families




**Figure 6:**

Average of 3 social deprivation indicators of residents in HPE compared to Ontario. The 3 indicators are: % of private households with only one person, % of people 15+ that are separated, divorced or widowed, and % of families that are lone parent families. Source: Census 2016

## PERSONAL HEALTH PRACTICES AND COPING SKILLS

Personal health practices or behaviours are the things one does in their daily life that can make them more or less healthy. This includes a variety of things such as the quantity of fruits and vegetables consumed, or the frequency of alcohol consumption. These activities performed every day can add up to have a large effect on health, influencing the risk of developing certain chronic illnesses.


### Health Behaviours of HPE Compared to Ontario



**Figure 7:** Health behaviours of residents in HPE compared to Ontario. Statistically significant differences are denoted by a star. Source: CCHS 2014

HPE has a significantly higher proportion of people that smoke and who are limited in their participation of physical activities when compared to Ontario. Relatedly, chronic conditions like arthritis, diabetes, asthma, mood disorders and COPD are also more prevalent in HPE. Smoking has been proven to be a cause of COPD.<sup>16</sup> Limitations to physical activity may be forcing more people to adopt a sedentary lifestyle, which will lead to higher rates of chronic illnesses like diabetes.<sup>17</sup>

### Health Outcomes of HPE Compared to Ontario



**Figure 8:** Health outcomes of residents in HPE compared to Ontario. Statistically significant differences are denoted by a star. Source: CCHS 2014



## HEALTHY CHILDHOOD DEVELOPMENT


The conditions and experiences that a child faces have been seen to carry implications throughout the rest of their adult life. For example, toxic stress, a prolonged negative stress that a child may face, will affect their development and may cause serious health issues. Stress and other negative circumstances are most damaging if experienced within the first 6 years of life, the most important time for brain development.<sup>8</sup> It is important to address such issues earlier, to prevent more serious downstream complications.

### EDI Vulnerability in HPE

The Early Development Instrument (EDI) score is a rating given to kindergarten students by their teachers. The students' vulnerability in terms of 5 different types of development is evaluated. The 5 categories are:

1. Physical health and well-being
2. Social competence
3. Emotional maturity
4. Language and cognitive development
5. Communication skills and general knowledge

#### % of children vulnerable in one or more sections of EDI



A child's development can be impacted by **negative circumstances** such as<sup>18-20</sup>:

- Poverty
- Parental divorce
- Abuse
- Institutionalization



Some **negative outcomes** associated with poor childhood development include<sup>21</sup>:

- Poor physical health
- Poor emotional health
- Lack of social competence
- Unhealthy coping behaviours



There are **protective factors** that may aid in childhood development including<sup>20</sup>:

- Emotional support from parents
- Positive relationships with adults outside family
- Support from organizations

**Figure 9:**

The EDI scores for HPE, by zone. Darker indicates a greater percentage of students that are vulnerable in one or more sections. Source: Hastings County Data Analysis Coordinator




# HEALTH SERVICES

Health service inaccessibility is a determinant of poor health. One is less likely to seek treatment for perceived health issues, legitimate or not, if they have low access to health services. In general, rural and less affluent areas suffer from lower number of health services, causing the population that live there to experience negative health outcomes.


In a recent study by the Institute for Clinical Evaluative Sciences (ICES), it was shown that there are differences in access to health services in Ontario based on where one lives.<sup>22</sup>

Large parts of HPE, like many other rural areas in southern Ontario, have very low accessibility to primary care physicians. Many of these places have a minimum driving distance of 30 minutes or more to the nearest physician, providing a substantial barrier to medical care.



**Figure 10:** Access to a comprehensive primary care physician within a 30 minute driving time in South Eastern Ontario 2014/2015. Source: ICES 2017

HPE has a high level of potentially avoidable mortality, which is the combination of both treatable and preventable causes of death when compared to Ontario. For both regions, avoidable mortality is a large portion of the premature mortality rate (Figure 11). This may be a consequence of low healthcare utilization, as lack of accessible health services is a major reason why people are not able to utilize health services. This is particularly true for the treatment of chronic conditions and for regular medical check-ups.<sup>23,24</sup>



Age-standardized rate (per 100 000 population)

**Figure 11:** Age-standardized mortality rates (per 100 000 people) for residents in HPE compared to Ontario. Source: CANSIM table 102 - 4315

## PHYSICAL ENVIRONMENT

The physical environment one lives in has many different effects on one's health. Some factors are more obvious such as water quality and air pollution, but other factors are more subtle such as the walkability of the area one lives in. Many places in HPE are physically isolated, and as a result many people commute by car. People traveling by car are not profiting from the health benefits associated with biking or walking, and are at a higher risk of cardiovascular disease.<sup>25</sup>

## SOCIAL ENVIRONMENT

The sense of community belonging one has and their engagement with their social environment can help support their health.<sup>26</sup> Being an active member of the local community can help someone cope with difficult circumstances. Being aware of available resources in the community may also serve to improve health. This determinant acts in ways similar to the social support network determinant of health. As such, someone with less personal support may benefit more from engaging in a positive social environment.

## CULTURE

Certain cultures are at a higher risk of health problems. For example, **Indigenous populations** suffer from lower income and lower levels of educational attainment.<sup>27</sup> This creates problems with food insecurity and crowded housing for Indigenous Canadians. The rates of infectious and chronic diseases are higher as well as suicide rates and substance abuse, when compared to the Canadian population.<sup>4</sup> This is of particular concern to HPE, where there is a larger aboriginal population compared to Ontario (5% vs. 2%) (NHS 2011).

**Immigrants** to Canada are another group that may be at a higher risk of negative health. Immigrants generally suffer from higher unemployment, underemployment, and social exclusion.<sup>4,28</sup> Conversely, there is a healthy immigrant effect which is when newly landed immigrants are actually found to be healthier than the Canadian population, but then eventually regress to Canadian levels.<sup>29</sup> There is a much smaller immigrant population in HPE compared to Ontario (7% vs. 29%) (NHS 2011).

## GENDER

Gender and the societal roles of gender play a part in determining one's health. Women tend to earn less, and be the parent in lone parent families more frequently than males,<sup>30</sup> putting them at higher risk of poverty and the negative health that accompanies it.<sup>25</sup> Males on the other hand are more susceptible to social exclusion and workplace related injuries, which bring about different health challenges.<sup>4</sup> Gender identity may also affect one's health; gay, lesbian and transgender youth are at a particularly high risk of developing mental health issues for example.<sup>31</sup>

### Written by:

Michael Elten, MSc Candidate, Foundational Standards Student  
Veronica Montgomery, MPH, Foundational Standards Manager

# REFERENCES

1. Statistics Canada. Health Indicators. 82-221-X., No. 1. Health Regions and Peer Groups. Ottawa, Ont: Minister of Industry; 2008. <http://www.statcan.gc.ca/pub/82-221-x/2008001/hregions-regionss/5202306-eng.htm>.
2. Halton Region. Attitudes and Beliefs about Social Determinants of Health. <http://www.halton.ca/cms/One.aspx?portalId=8310&pageId=120856>. Published May 2014, Accessed July 18, 2017.
3. Leeds, Grenville & Lanark District Health Unit. Attitudes & Beliefs about Social Determinants of Health in Leeds, Grenville & Lanark. [http://www.healthunit.org/reportpub/reports/Social\\_Determinants\\_of\\_Health-Attitudes\\_and\\_Beliefs.pdf](http://www.healthunit.org/reportpub/reports/Social_Determinants_of_Health-Attitudes_and_Beliefs.pdf). Published 2013. Accessed July 18, 2017.
4. Mikkonen J, Raphael D. *Social Determinants of Health - The Canadian Facts*. Toronto: York University School of Health Policy and Management. 2010:1-63.
5. Auger N, Alix C. Income, Income Distribution, and Health in Canada. In: Raphael D, ed. *Social Determinants of Health: Canadian Perspectives*. 2nd Edition. Toronto: Canadian Scholar's Press; 2009:90-107.
6. von Wagner C, Knight K, Steptoe A, Wardle J. Functional health literacy and health-promoting behaviour in a national sample of British adults. *J Epidemiol Community Health*. 2007;61(12):1086-1090. doi:10.1136/jech.2006.053967.
7. Blane D. Explanations of the difference in mortality risk between different educational groups. *Int J Epidemiol*. 2003;32:355-356. doi:10.1093/ije/dyg078.
8. Federal, Provincial and Territorial Advisory Committee on Population Health. *Toward a Healthy Future: Second Report on the Health of Canadians*. <http://publications.gc.ca/collections/Collection/H39-468-1999E.pdf>. Published September 1999. Accessed July 26 2017.
9. Uppal S. Young Men and Women without a High School Diploma. <http://www.statcan.gc.ca/pub/75-006-x/2017001/article/14824-eng.htm>. Published May 2017. Accessed August 4, 2017.
10. Pampalon R, Hamel D, Gamache P, Raymond G. A deprivation index for healthy planning. *Chronic Dis Can*. 2009;29(4):178-191.
11. Berkman LF, Syme SL. Social networks, host resistance, and mortality: a nine-year follow up study of Alameda County residents. *Am J Epidemiol*. 1979;109(2):186-204.
12. House J, Robbins C, Metzner H. The association of social relationships and activities with mortality: prospective evidence from the tecumseh community health study. *Am J Epidemiol*. 1982;116 (1):123-140.
13. Blazer DG. Social Support and Mortality in an Elderly Community Population. *Am J Epidemiol*. 1982;115(5):684-694. <http://aje.oxfordjournals.org/content/115/5/684>.
14. Berkman LF. Assessing the physical health effects of social networks and social support. *Annu Rev Public Health*. 1984;5:413-432.
15. Cohen S, Wills TA. Stress, Social Support, and the Buffering Hypothesis. *Psychol Bull*. 1985;98(2):310-357. doi:10.1037/0033-2909.98.2.310.
16. Lee PN, Forey BA, Coombs KJ. Systematic review with meta-analysis of the epidemiological evidence in the 1900s relating smoking to lung cancer. *BMC Cancer*. 2012;12:385. doi:10.1186/1471-2407-12-385.
17. Wilmot EG, Edwardson CL, Achana FA, et al. Sedentary time in adults and the association with diabetes, cardiovascular disease and death: Systematic review and meta-analysis. *Diabetologia*. 2012;55(11):2895-2905. doi:10.1007/s00125-012-2677-z.
18. Offord DR, Lipman EL. Lessening the impact of poverty on children. *Paediatr Child Heal*. 1999;4(8):526-529. [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=20212998](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20212998).
19. Mustard JF. Brain development, child development – Adult health and well-being and paediatrics. *Paediatr Child Heal*. 1999;4(8):519-520. [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=20212996](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20212996).
20. Maughan B, McCarthy G. Childhood adversities and psychosocial disorders. *Br Med Bull*. 1997;53(1):156-169. doi:10.1093/oxfordjournals.bmb.a011597.
21. Repetti RL, Taylor SE, Seeman TE. Risky Families: Family social environments and the mental and physical health of offspring. *Psychol Bull*. 2002;128(2):330-366.
22. Green ME, Gozdyra P, Frymire E, Glazier RH. Geographic Variation in the Supply and Distribution of Comprehensive Primary Care Physicians in Ontario. <https://www.ices.on.ca/Publications/Atlases-and-Reports/2017/Geographic-variation-in-physician-supply>. Published July 2017. Accessed July 18 2017.
23. Arcury TA, Gesler WM, Preisser JS, Sherman J, Spencer J, Perin J. The effects of geography and spatial behavior on health care utilization among the residents of a rural region. *Health Serv Res*. 2005;40(1):135-155. doi:10.1111/j.1475-6773.2005.00346.x.
24. Nemet GF, Bailey AJ. Distance and health care utilization among the rural elderly. *Soc Sci Med*. 2000;50(9):1197-1208. doi:10.1016/S0277-9536(99)00365-2.
25. Hamer M, Chida Y. Active commuting and cardiovascular risk: A meta-analytic review. *Prev Med*. 2007;46(1):9-13. doi:10.1016/j.ypmed.2007.03.006.
26. Zunzunegui M-V, Alvarado BE, Del Ser T, Otero A. Social networks, social integration, and social engagement determine cognitive decline in community-dwelling Spanish older adults. *J Gerontol B Psychol Sci Soc Sci*. 2003;58(2):S93-S100. doi:10.1093/geronb/58.2.S93.
27. Wilson D, Macdonald D. *The Income Gap between Aboriginal Peoples and the Rest of Canada*. Ottawa: Canadian Centre for Policy Alternatives; 2010 Apr.
28. Galarneau D, Morissette R. Immigrants: Settling for Less?. <http://www.statcan.gc.ca/pub/75-001-x/10604/6921-eng.pdf>. Published June 2004. Accessed Aug 4, 2017.
29. Deri C. Understanding the "Healthy Immigrant Effect" in Canada. <https://sciencessociales.uottawa.ca/economics/sites/socialsciences.uottawa.ca/economics/files/0502E.pdf>. Published January 2004. Accessed August 1, 2017.
30. Statistics Canada. 2016 Census of Canada: Families, households and marital status data table. <http://bit.ly/2xdQMpG>. Published August 2, 2017. Accessed August 3, 2017.
31. Marshal MP, Dietz LJ, Friedman MS, et al. Suicidality and depression disparities between sexual minority and heterosexual youth: A meta-analytic review. *J Adolesc Heal*. 2011;49(2):115-123. doi:10.1016/j.jadohealth.2011.02.005.