

Routine Publicly Funded Immunization Schedule for Ontario (Children Starting Immunization in Infancy*)

For more information, contact the Immunization Information line at 613-966-5500 x 221 www.hpepublichealth.ca

Immunizing Agent / OHIP Billing Code	2 Mos	4 Mos	6 Mos	12 Mos	15 Mos	18 Mos	4-6 Yrs	Gr 7-12	Gr 7-12	Gr 7-12	14-16 Yrs	Every 10 Yrs	65-70 Yrs	65+ Yrs
Diphtheria, tetanus, pertussis, polio, Hib Pediace / Pentace G841	✓1	✓1	✓1			✓1								
Pneumococcal conjugate 15 Vaxneuvance G846	✓2	✓2		✓2										
Rotavirus Rotarix (2 doses) N/A	✓3	✓3												
Meningococcal C conjugate Menjugate / NeisVac-C / Meningitec G844				✓4										
Measles, mumps, rubella (MMR) MMRII / Priorix G845				✓5										
Varicella Varivax III / Varilrix G848					✓6									
Measles, mumps, rubella, varicella (MMRV) Priorix-Tetra / Pro Quad G538							✓7							
Tetanus, diphtheria, pertussis, polio Adace-Polio / Boostrix-Polio G538							✓8							
Hepatitis B Recombivax HB / Engerix B G842								✓9						
Meningococcal conjugate ACYW-135 Menactra/ Nimenrix G538									✓10					
Human papillomavirus (HPV9) Gardasil G843										✓11				
Tetanus, diphtheria, pertussis Adace / Boostrix G847											✓12	✓12		
Tetanus, diphtheria Td absorbed G538												✓13		
Pneumococcal conjugate 20 Prevna 20 G846														✓14
Shingles Shingrix G538													✓15	
Influenza (seasonal) G590 Covid* G593			✓16											

*Based on Ministry Of Health recommendations

1	DTap-IPV-Hib Recommended at 2, 4, 6 and 18 months of age to complete the primary series.
2	Pneumo 15 Recommended at 2, 4, and 12 months of age.
3	Rotarix Recommended orally at 2 and 4 months of age. First dose should be given between 6 weeks of age and before 15 weeks of age; the second dose is given before 25 weeks of age. Doses must be at least 4 weeks apart.
4	Meningococcal-C Recommended at 12 months of age (or up to grade 6 if missed).
5	MMR Recommended on or after the first birthday; the second dose is Recommended at 4-6 years of age combined with varicella (MMRV).
6	Varicella Recommended at 15 months; the second dose is recommended between 4-6 years of age combined with MMR (MMRV).
7	MMRV Recommended for the second doses of MMR and varicella between 4-6 years of age. In Ontario, it is only recommended from 4-12 years of age inclusive.
8	Tetanus, diphtheria, pertussis, polio Recommended between 4-6 years of age to complete their primary childhood series.
9	Hepatitis B Recommended to all Grade 7-12 students under the School-Based Program. For those 11-15 years of age inclusive it is a 2 dose series and a 3-dose series for anyone outside this age range.
10	Meningococcal conjugate ACYW-135 Recommended for all Grade 7-12 students under the School-Based Program.
11	HPV9 Recommended to all Grade 7-12 students under the School-Based Program.
12	Tetanus, diphtheria, and pertussis Recommended for 14-16 year olds. Also publicly-funded as a one-time dose for adults who have never received an acellular pertussis-containing vaccine.
13	Tetanus, diphtheria Recommended every 10 years for adults. [Note: Tetanus, diphtheria, and pertussis is publicly funded as a one-time dose for adults, and should be given to those who have never received an acellular pertussis-containing vaccine.]
14	Pneumococcal conjugate 20: Recommended one time dose for individuals 65 years of age and older.
15	Shingles: Recommended for individuals 65-70 years of age inclusive.
16	Influenza: Recommended one dose every year in the fall. For ages 6 months and up Covid: Based on current MOH recommendations. For ages 6 months and up.

Additional doses may be needed for high-risk individuals, or for catch-up schedules. Refer to *Publicly Funded Immunization Schedules for Ontario* (2022 or as current) at ontario.ca/files/2024-01/moh-publicly-funded-immunization-schedule-en-2024-01-23.pdf

We are committed to providing accessible publications, programs and services to all. For assistance please call 613-966-5500; TTY: 711; or email accessibility@hpeph.ca. For more information, please visit hpepublichealth.ca.