

Publicly Funded Immunization Schedules for Ontario – January 2021

Publicly funded vaccines may be provided only to eligible individuals and must be free of charge

Routine Schedule: Children Starting Immunization in Infancy													
Vaccine	Age	2 Months	4 Months	6 Months	1 Year Φ	15 Months	18 Months	4 Years	Grade 7	14 Years	24 Years	≥ 34 Years Υ	65 Years
DTaP-IPV-Hib Diphtheria, Tetanus, Pertussis, Polio, <i>Haemophilus influenzae</i> type b		◆	◆	◆			◆						
Pneu-C-13 Pneumococcal Conjugate 13		◆	◆		◆								
Rot-5 Rotavirus		▲	▲	▲									
Men-C-C Meningococcal Conjugate C					◆								
MMR Measles, Mumps, Rubella					■								
Var Varicella						■							
MMRV Measles, Mumps, Rubella, Varicella								■					
Tdap-IPV Tetanus, diphtheria, pertussis, Polio								◆					
HB Hepatitis B									●				
Men-C-ACYW Meningococcal Conjugate ACYW-135									●				
HPV-9 Human Papillomavirus									●				
Tdap Tetanus, diphtheria, pertussis										◆	◆		
Td (booster) Tetanus, diphtheria												◆ Every 10 years	
HZ Herpes Zoster													I
Pneu-P-23 Pneumococcal Polysaccharide 23													■
Inf Influenza													
* Every year in the fall													
◆ - A single vaccine dose given by intramuscular injection ■ - A single vaccine dose given by subcutaneous injection ▲ - A single vaccine dose given by mouth ● - Provided through school-based immunization programs. Men-C-ACYW is a single dose. HB is a 2 dose series (see Table 6); HPV-9 is a 2 dose series (see Table 10). Each vaccine dose is given by intramuscular injection Φ - Given no earlier than the 1 st birthday, and prior to 16 months of age													
Υ - Once a dose of Tdap is given in adulthood (24 years of age), adults should receive Td boosters every 10 years thereafter I - HZ is a 2 dose series (see Table 12) given by intramuscular injection * - Children 6 months to 8 years of age who have not previously received a dose of influenza vaccine require 2 doses given ≥ 4 weeks apart. Children who have previously received ≥ 1 dose of influenza vaccine should receive 1 dose per season thereafter Note: A different schedule and/or additional doses may be needed for high risk individuals (see Table 3) or if doses of a vaccine series are missed (see appropriate Tables 4-24)													

◆ - Once a dose of Tdap is given in adulthood (24 years of age), adults should receive Td boosters every 10 years thereafter
■ - HZ is a 2 dose series (see Table 12) given by intramuscular injection
***** - Children 6 months to 8 years of age who have not previously received a dose of influenza vaccine require 2 doses given 24 weeks apart. Children who have previously received ≥1 dose of influenza vaccine should receive 1 dose per season thereafter
Note: A different schedule and/or additional doses may be needed for high risk individuals (see Table 3) or if doses of a vaccine series are missed (see appropriate Tables 4-24)

◆ - A single vaccine dose given by intramuscular injection
■ - A single vaccine dose given by subcutaneous injection
▲ - A single vaccine dose given by mouth
● - Provided through school-based immunization programs. Men-C-ACYW is a single dose; HB is a 2 dose series (see Table 6); HPV-9 is a 2 dose series (see Table 10). Each vaccine dose is given by intramuscular injection
Φ - Given no earlier than the 1st birthday, and prior to 16 months of age

Publicly Funded Immunization Schedules for Ontario – January 2021

Publicly funded vaccines may be provided only to eligible individuals and must be free of charge

Catch-up Schedule 1: Children Starting Immunization between 1-6 Years																			
Age Vaccine		1 st Visit:			2 nd Visit: 2 months after 1 st visit				3 rd Visit: 2 months after 2 nd visit		4 th Visit: 6-12 months after 3 rd visit		5 th Visit (only required if child was <4 years at 4 th visit) 4-6 yrs of age and 6-12 months after 4 th visit	Grade 7	14-18 yrs§	24-28 yrs		≥34, yrs†	65 yrs
		If child is			If child is <5 years and was				If child is		If child is		If child is				If adult was		
		<4 yrs	4 yrs	5-6 yrs	<2 yrs at 1 st visit	2-3 yrs at 1 st visit	4 yrs at 1 st visit	5-6 yrs	7 yrs	<7 yrs	7 yrs	<4 yrs	4-8 yrs			<18 yrs at previous visit†	≥18 yrs at previous visit†		
DTaP-IPV-Hib		◆	◆	◆	◆	◆	◆					◆							
Pneu-C-13		◆	◆		◆														
MMR		■																	
MMRV			■	■								■	■						
Var					■	■													
Men-C-C		◆	◆	◆															
Tdap-IPV									◆				◆	◆					
HB														●					
Men-C-ACYW														●					
HPV-9														●					
Tdap																	◆		
Td																◆	◆	◆ Every 10 years	
HZ																			■
Pneu-P-23																			■
Inf																			
* Every year in the fall																			
◆ - A single vaccine dose given by intramuscular injection																			
■ - A single vaccine dose given by subcutaneous injection																			
● - Provided through school-based immunization programs. Men-C-ACYW is a single dose; HB is a 2 dose series (see Table 6); HPV-9 is a 2 dose series (see Table 10). Each vaccine dose is given by intramuscular injection																			
§ - Given 10 years after the (4-8 year old) Tdap-IPV dose																			
† - Given 10 years after the adolescent Tdap dose																			
* Every year in the fall																			
† - Once a dose of Tdap is given in adulthood (>18 yrs), adults should receive Td boosters every 10 years thereafter																			
■ - HZ is a 2 dose series (see Table 12) given by intramuscular injection																			
★ - Children 6 months to 8 years of age who have not previously received a dose of influenza vaccine require 2 doses given 24 weeks apart. Children who have previously received ≥1 dose of influenza vaccine should receive 1 dose per season thereafter																			
Note: A different schedule and/or additional doses may be needed for high risk individuals (see Table 3) or if doses of a vaccine series are missed (see appropriate Tables 4-24)																			

Publicly Funded Immunization Schedules for Ontario – January 2021

Publicly funded vaccines may be provided only to eligible individuals and must be free of charge

Catch-up Schedule 2: Children Starting Immunization between 7-17 Years														
Age Vaccine	1 st Visit		2 nd Visit: 2 months after 1 st Visit		3 rd Visit: 6-12 months after 2 nd Visit	Grades 7-8	Grades 7-12	Grades 7-12 Males born in or after 2004	Grades 7-12 Females	10 Years after 3 rd Visit	10 years after previous visit (only required if child was <18 yrs old at previous visit)	Every 10 years after the previous visit ^a	65 Years	
	If child is <13 yrs	If child is 13 to 17 yrs	If child is <13 yrs	If child is 13 to 18 yrs										
Tdap-IPV	◆	◆	◆	◆	◆									
MMRV	■		■											
MMR		■		■										
Var		■		■										
Men-C-C	●													
HB						●								
Men-C-ACYW							●							
HPV-9								●						
Tdap									●	◆				
Td												◆		
HZ													■	
Pneu-P-23													■	
Inf														

◆ - A single vaccine dose given by intramuscular injection

■ - A single vaccine dose given by subcutaneous injection

● - Individuals born on or after 2003/Sept/01 are eligible to receive a dose of Men-C-C (given by intramuscular injection). These individuals are also eligible to receive Men-C-ACYW when they enter Grade 7. If the individual is immunized with Men-C-ACYW, in or after Grade 7, Men-C-C is no longer recommended

● - Provided through school-based immunization programs. Men-C-ACYW is a single dose; HB is a 2 dose series (see Table 6); HPV-9 is a 2 or 3 dose series (see Tables 10 and 11). Each vaccine dose is given by intramuscular injection

◆ - Once a dose of Tdap is given in adulthood, adults should receive Td boosters every 10 years thereafter

■ - HZ is a 2 dose series (see Table 12) given by intramuscular injection

◆ - Children 6 months to 8 years of age who have not previously received a dose of influenza vaccine require 2 doses given 24 weeks apart. Children who have previously received ≥1 dose of influenza vaccine should receive 1 dose per season thereafter

Note: A different schedule and/or additional doses may be needed for high risk individuals (see Table 3) or if doses of a vaccine series are missed (see appropriate Tables 4-24)

Catch-up Schedule 3: Adults Starting Immunization at 18 Years and Older										
Age Vaccine	1 st Visit			2 nd Visit: 2 months after 1 st Visit				3 rd Visit: 6-12 months after 2 nd Visit	Every 10 years after the 3 rd Visit	65 Years
	In or prior to 1985	between 1986 and 1996	If adult is born between 1997 and 1999	In or after 2000	In or after 2000 and is 19 to 25 yrs	In or prior to 1999 and is 20 to 25 yrs	If adult is born in or prior to 1999 and is ≥26 yrs			
Tdap-IPV	◆	◆	◆	◆						
MMR	■	■	■	■	■	■				
Var				■	■					
Men-C-ACYW			◆	◆						
Men-C-C		◆								
Td					◆	◆	◆	◆	◆	
IPV					◆	◆		◆		
HZ										■
Pneu-P-23										■
Inf										
* Every year in the fall										
◆ - A single vaccine dose given in a syringe and needle by intramuscular injection ■ - A single vaccine dose given in a syringe and needle by subcutaneous injection ■ - HZ is a 2 dose series (see Table 12) given by intramuscular injection Note: A different schedule and/or additional doses may be needed for high risk individuals (see Table 3) or if doses of a vaccine series are missed (see appropriate Tables 4-24)										